


Abstract

Background: The 3D structure and function of proteins is influenced by post-translational modifications. Palmitoylation, one such modification, is the covalent attachment of a 16-carbon fatty acid chain to cysteine residues. Palmitoylation facilitates protein association with membranes due to increased hydrophobicity. Unlike other lipid modifications in which the protein is permanently modified, protein palmitoylation is dynamic, and palmitate groups are added and removed as the cell requires. Palmitate groups are attached to proteins by protein acyl transferases, and removed by protein thioesterases, such as palmitoyl protein thioesterase 1 (PPT1), which breaks the thioester link between the palmitate and the protein. In neurons, PPT1 is enriched at synapses, and its dysfunction leads to aberrant increases in palmitoylation of synaptic proteins such as SNAP-25, cysteine string protein α , and synaptobrevin 2. Mutations in the PPT1 gene (CLN1) lead to autosomal recessive infantile neuronal ceroid lipofuscinosis (INCL), a progressive neurodegenerative disease. CLN1 mutations lead to a loss of the depalmitoylation activity of PPT1 and result in synaptic trafficking deficits. However, the repertoire of substrates of PPT1 is unknown. **Aim:** To identify synaptic PPT1 substrates. **Method:** We purified palmitoylated proteins from wild-type and PPT1 knockout (KO) synaptosomes and compared the palmitomes using Label Free Quantification-Mass Spectroscopy. **Results:** We identified putative PPT1 substrates and will validate select synaptic substrates using orthologous methods. We also mapped and characterized the synaptic pathways most affected by PPT1 KO. **Conclusion:** Our results reveal the critical roles PPT1 plays in synapse function and protein depalmitoylation deficits in CLN1.

Palmitoyl Protein Thioesterase 1 (PPT1)


Palmitoylation is a dynamic post-translational modification that entails the covalent attachment of a 16-carbon fatty acid chain typically to cysteine residues. Palmitate groups are added by palmitoyl transferases such as the DHHC proteins and are removed by thioesterases such as PPT1, which cleave the thioester linkage. Palmitoylation allows proteins to associate with membranes, however depalmitoylation may be required for lysosomal degradation.




Neuronal ceroid lipofuscinoses

Neuronal ceroid lipofuscinoses are a family of hereditary lysosomal storage disorders characterized by the accumulation of the autofluorescent pigment lipofuscin in the lysosomes. These lysosomal accumulations also contain highly lipidated peptides.

Loss-of function mutations in PPT1 are associated with infantile NCL, an autosomal recessive form with rapid progression. The average lifespan of INCL patients is 9-11 years.




Isolation of the synaptic palmitome



Scheme for identification of PPT1 substrates. (A) WT and PPT1 KO synaptosomes were assessed to determine baseline protein levels (proteome) and palmitoylated protein levels (palmitome) following Acyl Resin-Assisted Capture (Acyl-RAC). (B) Acyl-RAC is a method to isolate palmitoylated proteins on thiopropyl sepharose beads. Levels of isolated proteins can be quantified to assess palmitome changes in PPT1 KO vs. WT.

PPT1 KO results in changes to the synaptic palmitome



Accession	Description	KO/WT	p-value
LRTM2*	Leucine-rich repeat and transmembrane domain-containing protein 2	18.450	0.0273
SATT*	Neutral amino acid transporter A	8.375	0.0440
ITM2B*	Integral membrane protein 2B	6.760	0.0071
VPS29	Vacuolar protein sorting-associated protein 29	6.603	0.0403
AT1B2*	Sodium/potassium-transporting ATPase subunit beta-2	5.414	0.0177
GBRG2*	Gamma-aminobutyric acid receptor subunit gamma-2	5.310	0.0353
GPM6A*	Neuronal membrane glycoprotein M6-a	5.105	0.0068
CATD	Cathepsin D	4.751	0.0038
RMD3*	Regulator of microtubule dynamics protein 3	4.494	0.0453
DNJC5	DnaJ homolog subfamily C member 5	4.358	0.0109
SYUA	Alpha-synuclein	4.021	0.0497
VAMP1*	Vesicle-associated membrane protein 1	4.005	0.0172

Table 1. Subset of the 214 proteins identified as putative PPT1 substrates that will be validated using orthologous methods. Once validated, we will assess localization changes of PPT1 substrates without transmembrane domains (transmembrane domain indicated by asterisk) in PPT1 KO vs WT neurons by fractionation. Notably, proteins involved in GABA receptor signaling (GBRG2) are identified as putative PPT1 substrates. GABA receptor signaling was identified by Ingenuity Pathway analysis as the most affected pathway in PPT1 KO vs WT palmitome samples.

* indicates known transmembrane domain.

Future Directions

We plan to validate hits from Acyl-RAC analysis by incubation of synaptosomes with purified PPT1 followed by Acyl-RAC to assess whether PPT1 itself is able to depalmitoylate putative substrates. As palmitoylation is an activity-dependent modification, we will also stimulate neurons prior to Acyl RAC to identify substrates whose depalmitoylation by PPT1 is activity dependent.

We also plan to screen PPT1 substrates for consensus sequences that may be recognized by PPT1.

Finally, we would like to assess how loss of PPT1 function results in neurodegenerative disease by studying NCL patient neurons and assessing substrate palmitoylation.

Acknowledgements

We would like to thank Wilhemina Koomson for initial experiments. This work was funded by: NIH 1R01 NS083846, NIH 1R21 NS094971, the Yale Gruber science fellowship and the NIDA Neuroproteomic center NIH DA018343.

